

THE b -CHROMATIC NUMBER IN REGULAR GRAPHS

MAGDA DETTLAFF, HANNA FURMAŃCZYK, RADOSŁAW ZIEMANN

University of Gdańsk

e-mail: magda.dettlaff@ug.edu.pl, hanna.furmanczyk@ug.edu.pl,
radoslaw.ziemann@ug.edu.pl

A b -coloring of the vertices of a graph G is a proper coloring where each color class contains a vertex which is adjacent to a vertex in each other color class. The b -chromatic number of G is the maximum integer $\chi_b(G)$ for which G has a b -coloring with $\chi_b(G)$ colors. In our poster we give some partial results for one of the most important conjectures regarding the b -chromatic number of regular graphs which was first posted by El Sahili and Kouider in 2009:

Every d -regular graph of girth at least 5, different than the Petersen graph, has a b -coloring with $d + 1$ colors.

References

- [1] A. El Sahili, H. Kouider, About b -colouring of regular graphs. Util. Math. 80 (2009), 211–215.
- [2] M. Blidia, F. Maffray, Z. Zemir, On b -colorings in regular graphs. Discrete Appl. Math. 157 (2009), 1787–1793.
- [3] M. Jakovac, I. Peterin, The b -chromatic number and related topics—A survey. Discrete Appl. Math. 235 (2018), 184–201.